



**UNIVERSITI PUTRA MALAYSIA**

**ESTABLISHMENT OF TISSUE CULTURE AND HAIRY ROOT  
PRODUCTION SYSTEMS FOR SOLENOSTEMON  
SCUTELLARIOIDES (L.) CODD**

**LE VINH THUC.**

**FBSB 2004 6**

**ESTABLISHMENT OF TISSUE CULTURE AND HAIRY ROOT  
PRODUCTION SYSTEMS FOR  
*Solenostemon scutellarioides* (L.) Codd**

**By**

**LE VINH THUC**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the  
Degree of Master of Science**

**September 2004**



**Dedicated to my parents**

**Mr. Lê Văn Khoản and Mrs. Trần Thị Kim Đăng**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the degree of Master of Science

**ESTABLISHMENT OF TISSUE CULTURES AND HAIRY ROOT  
PRODUCTION SYSTEMS FOR  
*Solenostemon scutellarioides* (L.) Codd**

**By**

**LE VINH THUC**

**September 2004**

**Chairman: Associate Professor Norihan Mohd. Saleh, Ph.D.**

**Faculty: Biotechnology and Biomolecular Sciences**

*In vitro* propagation of *S. scutellarioides* was carried out by culturing shoot tip explants onto LS, MS and B5 basal medium supplemented with different concentrations of BAP. Comparative analysis of the different types of basal media showed that the LS basal medium supplemented with 8.88  $\mu\text{M}$  BAP was suitable for multiple shoots formation on *S. scutellarioides* (~ 13 shoots/explant). Addition of higher concentrations of BAP (35.5  $\mu\text{M}$ ) inhibited multiple shoots and induced abnormal plantlet formation in this species.

Culturing shoot tip explants on B5 medium supplemented with 4.52  $\mu\text{M}$  BAP and 2.26  $\mu\text{M}$  2,4-D produced pink friable callus. On the other hand, culturing leaf explants on MS and B5 media both supplemented with 4.52  $\mu\text{M}$  2,4-D and 0.46  $\mu\text{M}$  2,4-D, produced yellowish and grayish friable calli, respectively. Following calli induction, cell suspension culture of *S. scutellarioides* was successfully initiated on MS medium supplemented with 2.26  $\mu\text{M}$  2,4-D and 0.47  $\mu\text{M}$  kinetin. The cell suspension required only half the amount of 2,4-D

required for calli induction on solid medium (MS + 4.52  $\mu$ M 2,4-D + 0.46  $\mu$ M kinetin).

In an effort towards achieving genetically stable plant tissue culture material, for the future induction of valuable secondary metabolite, adventitious roots and hairy roots were induced from *S. scutellarioides* explants. Quantitative and qualitative assessments of the biomass producing from each of the culture method were analyzed.

Adventitious root cultures in *S. scutellarioides* leaves were induced by placing explants onto MS basal medium supplemented with different concentrations of auxins (NAA, IBA, IAA) and cytokinin (kinetin). Comparative analysis of the roots enhanced showed that MS medium supplemented with 5.0  $\mu$ M IBA produced rapidly growing adventitious roots. The presence of cytokinin (kinetin) was inhibiting to adventitious root formation in *S. scutellarioides*. Supplementing MS medium with auxin either IBA or NAA was enough to induce adventitious root formation.

Hairy roots of *S. scutellarioides* were induced by inoculation of leaf explants with *A. rhizogenes* strains TR 105, LBA 9402, 8196 and ATCC 15834. These strains showed different abilities to induce hairy root formation in the leaf explants. Assessment of the plant susceptibility to the different *A. rhizogenes* strains showed that the strains ATCC 15834, TR 105, LBA 9402, and 8196 produced 56.3%, 25.5%, 21.5%, and 13.8% transformation efficiencies, respectively. Acetosyringone was found to be useful for enhancement of hairy root production in *S. scutellarioides*.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

**PEMBANGUNAN TISU KULTUR DAN SISTEM PRODUKTIVITI AKAR  
RERAMBUT TUMBUHAN**

***Solenostemon scutellarioides* (L.) Codd**

Oleh

**LE VINH THUC**

**September 2004**

**Pengerusi: Profesor Madya Norihan Mohd Saleh, Ph.D.**

**Fakulti: Bioteknologi dan Sains Biomolekul**

Pembiakan secara *in vitro* dilakukan keatas eksplan pucuk *S. scutellarioides* menggunakan media LS, MS dan B5 yang mengandungi kepekatan BAP berbeza. Perbandingan ke atas media yang digunakan menunjukkan pertumbuhan pucuk berganda *S. scutellarioides* (~ 13 pucuk/eksplan) adalah paling sesuai dalam basal medium LS yang mengandungi 8.88  $\mu\text{M}$  BAP. Penambahan kepekatan BAP yang tinggi (35.5  $\mu\text{M}$ ) memberi kesan rencatan terhadap perkembangan pucuk berganda bagi tumbuhan ini.

Pengkulturan eksplan pucuk diatas medium B5 yang mengandungi 4.52  $\mu\text{M}$  BAP dan 2.26  $\mu\text{M}$  2,4-D menghasilkan kalus berwarna merah muda yang mudah terlerai. Manakala eksplan daun menghasilkan kalus kuning dan mudah terurai diatas MS yang mengandungi 4.52  $\mu\text{M}$  2,4-D dan 0.46  $\mu\text{M}$  serta kalus kelabu yang mudah terlerai di atas medium B5 dengan kandungan hormon yang sama. Seterusnya ampaian sel *S. scutellarioides* telah berjaya diinisiasikan dalam medium MS yang mengandungi

2.26  $\mu\text{M}$  2,4-D dan 0.46  $\mu\text{M}$  kinetin. Ampaian sel ini memerlukan hanya separuh sahaja kandungan 2,4-D berbanding dengan medium pepejal bagi mengaruh kalus (MS +  $\mu\text{M}$  2,4-D + 0.46  $\mu\text{M}$  kinetin).

Dalam usaha ke arah memperoehi kestabilan genetik kultur tumbuhan bagi tujuan menghasilkan metabolit sekunder yang bernilai dimasa depan, akar adventitious dan akar rerambut juga telah diaruh daripada eksplan *S. scutellarioides*.

Penghasilan biomass daripada kaedah kultur akar adventitious dan rerambut dinilai secara kuantitatif dan kualitatif ke atas biomass yang dihasilkan menggunakan kedua-dua kaedah pengkulturan telah dilakukan. Akar adventitious *S. scutellarioides* telah diaruh daripada eskplan daun menggunakan medium MS yang mengandungi kepekatan auksin (NAA, IBA, IAA) dan sitokinin (kinetin) berbeza. Analisia menunjukkan medium MS yang mengandungi kepekatan 5.0  $\mu\text{M}$  IBA menghasilkan pertumbuhan akar adventitious paling cepat. Terdapat kesan kerencatan akar adventitious *S. scutellarioides* dengan kehadiran sitokinin (kinetin). Perambahan auksin (IBA, NAA) dalam medium MS sahaja sudah memadai untuk mengaruh penghasilan akar adventitious.

Akar rerambut telah diaruh daripada eskplan daun *S. scutellarioides* dengan menggunakan strain TR105, LBA 9402, 8196 dan ATCC 15834 *A. rhizogenes*. Kesemua strain ini menunjukkan keupayaan berbeza dalam pembentukan akar rerambut. Penilaian sensitiviti tumbuhan ini terhadap

strain *A. rhizogenes* tersebut menunjukkan keberkesanan transformasi 56.3% bagi ATCC 15834, 25,5% bagi TR 105, 21.5% bagi LBA 9402 dan 13.8% bagi 8196 adalah berbeza. Kehadiran asetosringone penting untuk meningkatkan keberkesanan aruhan akar rerambut bagi *S. scutellarioides*.



## **ACKNOWLEDEMENTS**

First of all, I would like to express my most sincere gratitude and deepest appreciation to Assoc. Prof. Dr. Norihan Mohd. Saleh, my supervisor, Head of Cell and Molecular Biology Department, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Malaysia, for giving me the opportunity to pursue my studies in the field of plant biotechnology. With her constant motivation and words of advice, has inspired me to become a better researcher and individual. Special thanks are extended to other members of my committee, Prof. Abdul Manaf Ali and Dr. Mohd. Azmuddin Bin Abdullah for their advice, comments and guidance whenever sought.

I would like to thank Cantho University, Vietnam by giving me scholarship to study in Universiti Putra Malaysia, Malaysia for two years. Deep gratitude is also acknowledged to my Vietnamese lecturers and colleagues in Crop Science Department, College of Agriculture, Cantho University, Vietnam with their motivations helping me to be eager to study in Malaysia.

Sincere thanks also go to the staff in Tissue Culture Laboratory, Genetic Laboratory and Fitotron Laboratory and the official staff in Department of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Malaysia for their enthusiastic helps in laboratorial materials and in paper works, respectively.



Last but not least, I wish to express my deepest gratitude and appreciation to my family for their constant support throughout my study. To my best friend Kak Norrizah thank you for your help and criticism of my work. I also would like to thank all my friends while I studied here.

I certify that an Examination Committee met on 2<sup>nd</sup> September 2004 to conduct the final examination of Le Vinh Thuc on his Master of Science thesis entitled "Establishment of Tissue Culture and Hairy Root Production Systems for *Solenostemon scutellarioides* (L.) Codd" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**RAHA ABDUL RAHIM, Ph.D.**

Associate Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Chairman)

**SUHAIMI NAPIS, Ph.D.**

Associate Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Member)

**SITI NOR AKMAR ABDULLAH, Ph.D.**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**ISMANIZAN ISMAIL, Ph.D.**

Associate Professor  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
(Independent Examiner)



**GULAM/RUSUL RAHMAT ALI, Ph.D.**  
Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 21 SEP 2004

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**NORIHAN MOHD. SALEH, Ph.D.**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairperson)

**ABDUL MANAF ALI, Ph.D.**

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

**MOHD. AZMUDDIN BIN ABDULLAH, Ph.D.**

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)



**AINI IDERIS, Ph.D.**

Professor/Dean

School of Graduate Studies

Universiti Putra Malaysia

Date : 25 OCT 2004

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

  
LE VINH THUC

Date: 20 Sept. 2004

## TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xix
 <b>CHAPTER</b>	
 <b>1      INTRODUCTION</b>	 <b>1</b>
 <b>2      LITERATURE REVIEW</b>	 <b>5</b>
2.1 <i>Solanostemon scutellarioides</i>	5
2.1.1 Plant Description and Distribution	5
2.1.2 Secondary Metabolite Compounds from <i>S. scutellarioides</i>	7
2.2 <i>In vitro</i> Culture of <i>S. scutellarioides</i>	8
2.3 Tissue Culture Media	9
2.3.1 Choice of Basal Media	9
2.3.2 Vitamins	13
2.4 Plant Growth Regulators in Plant Tissue Cultures	14
2.4.1 Auxins	16
2.4.2 Cytokinins	17
2.5 Callus and Cell Suspension cultures	19
2.5.1 Callus Induction	19
2.5.2 Cell Suspension Culture	20
2.6 Organ Cultures	22
2.6.1 Shoot Culture	22
2.6.2 Normal Root Culture	23
2.6.3 Hairy Roots	27
2.7 Elicitation of Secondary Metabolites	32
 <b>3      IN VITRO PROPAGATION OF S. SCUTELLARIOIDES</b>	 <b>35</b>
3.1 Introduction	35
3.2 Materials and Methods	36
3.2.1 Plant Materials	36
3.2.2 Explant Sterilization	36
3.2.3 Culture Maintenance	37
3.2.4 Media Preparation	37
3.2.5 <i>In vitro</i> Propagation	38
3.2.6 Data Collection	38



3.2.7 Statistical Analysis	39
3.3 Results and Discussion	39
3.3.1 Explant Sterilization	39
3.3.2 Effect of Different Basal Media	41
3.3.3 Effect of BAP on the Growth of Explant	43
3.3.4 Effect of Different Basal Media and BAP on Multiple Shoot	47
3.4 Conclusion	50
<b>4 CALLUS INDUCTION AND INITIATION OF CELL SUSPENSION CULTURES OF <i>S. SCUTELLARIOIDES</i></b>	<b>52</b>
4.1 Introduction	52
4.2 Materials and Methods	53
4.2.1 Plant Materials	53
4.2.2 Media Preparation	54
4.2.3 Callus Induction from Various Explants	54
4.2.3.1 Shoot Tip Explants	54
4.2.3.2 Leaf Explants	55
4.2.4 Cell Suspension Cultures	55
4.2.5 Statistic Analysis	56
4.3 Results and Discussion	56
4.3.1 Callus Formation	56
4.3.1.1 Shoot Tip Explants	56
4.3.1.2 Leaf Explants	60
4.3.2 Initiation of Cell Suspension Cultures	64
4.4 Conclusion	68
<b>5 HAIRY ROOT AND ROOT CULTURES OF <i>S. SCUTELLARIOIDES</i></b>	<b>69</b>
5.1 Introduction	69
5.2 Materials and Methods	70
5.2.1 Plant Materials	70
5.2.2 Bacterial Strains	70
5.2.3 Activation of Bacterial Strains	72
5.2.4 Media Preparation	72
5.2.5 Hairy Root Cultures	73
5.2.5.1 Induction	73
5.2.5.2 Maintenance Hairy Roots	74
5.2.6 Adventitious Root Induction	74
5.2.6.1 Induction	74
5.2.6.2 Proliferation	75
5.2.6.3 Root Suspension	75
5.2.7 Data collection and Statistical Analysis	76
5.2.7.1 Hairy Root Experiment	76
5.2.7.2 Adventitious Root Induction Experiment	76
5.2.7.3 Statistical Analysis	76
5.3 Results and Discussion	77
5.3.1 Effect of Different <i>A. rhizogenes</i> Strains on Hairy Root Induction	77

5.3.2 Effect of Plant Growth Regulators on Adventitious Root Induction	81
5.3.3 Adventitious Roots Growing in Solid Media	90
5.3.4 Root Growing in Liquid Media	93
5.4 Conclusion	96
<b>6 GENERAL DISCUSSION</b>	<b>98</b>
BIBLIOGRAPHY	108
APPENDICES	131
BIODATA OF THE AUTHOR	140
PUBLISHED ACTIVITIES FROM THE THESIS	141



## LIST OF TABLES

Table	Page
2.1 Components of medium formulations	11
2.2 Shoot cultures of medicinal plants	24
2.3 Root cultures of medicinal plants	26
2.4 Hairy root cultures producing pharmaceutical products of interest	31
3.1 Effect of sterilization time on the percentage of contamination and death of <i>S. scutellarioides</i> explants	40
3.2 Effect of MS, LS, and B5 basal medium on percentage of increment in height and in fresh weight of explants in 2 months of culture	42
3.3 Percentage of increment in height and fresh weight of explants on different media in 2 months of culture	44
4.1 Quantity and quality assessment of 2 months old callus induced from shoot tips of <i>S. scutellarioides</i>	58
4.2 Callus induction on MS and B5 basal medium supplemented with various combinations of auxin and cytokinin in 2 months of culture	61
4.3 Effect of various concentrations of plant growth regulators on percentage of increment in fresh weight of suspended cells of <i>S. scutellarioides</i> in 1 month of culture	66
5.1 Characteristic of <i>A. rhizogenes</i> strains TR 105, LBA 9402, 8196, and ATCC 15834	71
5.2 The occurrence of hairy root induction (number of days in culture and its frequencies) in <i>S. scutellarioides</i> leaf explants in the presence or absence of acetosyringone induced by different strains of <i>A. rhizogenes</i>	78
5.3 Adventitious root formation on MS medium supplemented with various concentrations of auxin and cytokinin	85



5.4	Fresh weight and characteristic of adventitious roots on MS medium supplemented with different concentrations of auxin and cytokinin in 1 month old culture	87
5.5	The percentage of increment in fresh weight and characteristics of adventitious root on MS solid medium supplemented with auxins in 1 month old culture	91
5.6	Percentage of increment in fresh weight and characteristics of roots in MS liquid medium supplemented with various concentrations of auxins in in 1 month old culture	94

## LIST OF FIGURES

Figure	Page
2.1 <i>Solanostemon scutellarioides</i> in the field	6
2.2 Schematic diagram of the <i>Agrobacterium</i> infection process. Critical steps that occur to or within the bacterium (chemical signaling, <i>vir</i> gen induction, and T-DNA processing) and within the plant cell (bacterial attachment, T-DNA transfer, nuclear targeting, and T-DNA integration) are highlighted, along with genes and/or proteins known to mediate these events (Gelvin, 2000)	30
3.1 Characteristic of plantlet: A) big leaves, normal and roots (+); B) small leaves, normal and few roots (++); C) small leaves, few roots and abnormal (+++).	45
3.2 Addition of different concentrations of BAP in basal media (MS, LS, B5) on number of multiple shoot formation	49
4.1 Callus formation on B5 basal medium supplemented with 4.52 $\mu$ M BAP and 2.26 $\mu$ M 2,4-D in 2 months of culture.	59
4.2 Standard callus/explant used for scoring the callus induction: A) +++; B) ++; C) +; Arrows indicate not callus induced from explant	62
4.3 Cell suspension of <i>S. scutellarioides</i> : A) cultured on MS basal medium supplemented with 2.26 $\mu$ M of 2,4-D and 0.46 $\mu$ M of kinetin, B) cultured on B5 basal medium supplemented with 4.52 $\mu$ M of 2,4-D and 0.46 $\mu$ M of kinetin; black arrow indicate dead cell; red arrow indicate fine cells; blue arrow indicate clumpy cells	67
5.1 Hairy roots appeared at the cut edge of a leaf explant of <i>S. scutellarioides</i> . Arrows indicate the hairy root induced on different parts of the explant	82
5.2 Hairy roots of <i>S. scutellarioides</i> growing on hormone-free MS medium	83
5.3 Characteristics of adventitious root: A) normal; B) slightly swollen; C) swollen; Arrows indicate the swollen root	88

5.4	Characteristics of proliferated root: A) normal; B) slightly swollen; C) swollen; Arrows indicate swollen roots.	92
5.5	Characteristics of root in suspension culture: A) normal; B) succulent.	95

## LIST OF ABBREVIATION

%	Percentage
μM	Micromolar
2,4-D	2,4-dichlorophenoxyacetic acid
2iP	N6-isopentyladenine
<i>A. rhizogenes</i>	<i>Agrobacterium rhizogenes</i>
ANOVA	Analysis of variance
Auxin	Plant growth regulator resembling IAA in physiological activity
B5	Gamborg medium
BAP	6-Benzylaminopurine
cm	Centimeter
CPA	p-chlorophenoxyacetic acid
Cytokinin	Plant growth regulator stimulating cell division and resembling kinetin in physiological activity. Mainly N <sub>6</sub> substituted aminopurine compounds.
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleotic acid
EDTA	Ethylenediaminetetraacetic acid
Explant	Excised fragment of plant tissue or organ used to initiate a tissue culture
μg	Microgram
IAA	Indole-3-Acetic Acid
IBA	Indole-3-Butyric Acid
Kinetin	6-Furfurylaminopurine
l	Liter
LS	Linsmaier and Skoog
ml	milliliter
mm	Millimeter

<b>MS</b>	<b>Murashige and Skoog</b>
<b>MW</b>	<b>Molecular weight</b>
<b>NAA</b>	<b><math>\alpha</math>-Naphthalene acetic acid</b>
<b>°C</b>	<b>Degree centigrade</b>
<b>OD</b>	<b>Optical density</b>
<b>Ri</b>	<b>Root inducing</b>
<b>rpm</b>	<b>Revolution per minute</b>
<b><i>S. scutellarioides</i></b>	<b><i>Solanostemon scutellarioides</i></b>
<b>T-DNA</b>	<b>Transferable-DNA</b>
<b>T<sub>L</sub></b>	<b>Left border of T-DNA</b>
<b>T<sub>R</sub></b>	<b>Right border of T-DNA</b>
<b><i>Vir</i></b>	<b>Virulence</b>

## CHAPTER 1

### INTRODUCTION

Plant secondary metabolites are known to play a major role in the adaptation of plants to their environment. They are also important sources of active pharmaceuticals, food additives, cosmetic industries and pesticides such as rosmarinic acid, vinblastine and ginsenoside. However, useful secondary metabolites distributed in plants are rather more restricted than primary metabolites such as nucleotides, nucleic and amino acids, protein and carbohydrates. In addition, secondary metabolites are found in a few species or even within a few varieties within a species (Alfermann and Petersen, 1995). They are also limited in occurrence in the ornamental plants (Ramawat, 1999a). Rosmarinic acid for example occurs especially within the family of Boraginaceae and the subfamily Nepetoideae within the Labiate (Pedersen, 2000). They mostly accumulate in plant in small quantities, sometimes in specialized cells, making their extraction from the plant often difficult (Sasson, 1991).

Several studies suggested that production of plant secondary metabolites can be improved by using tissue and cell culture techniques (Bourgaud *et al.*, 2001; Suresh *et al.*, 2001; Luczkiewicz *et al.*, 2002). Cell suspension cultures of *Coleus blumei* for example are known to produce rosmarinic acid (Razzaque and Ellis, 1977). Sixty fold increases in anthraquinones

production from 0.3% dry weight to 18% dry weight have been reported in the cell suspension cultures of *Morinda citrifolia* (Misawa, 1994). *In vitro* culture systems such as cell suspension, callus, root, shoot and the transformed 'hairy root' cultures (Scragg, 1995) have been extensively studied with the objective of improving the production of secondary plant compounds (Gontier *et al.*, 1994; Bourgaud *et al.*, 2001).

Plant cell and organ cultures may provide steady and controlled environment for long-term supplies of plant secondary metabolites and protein (Bhadra *et al.*, 1993), enhanced production of pharmaceutically valuable alkaloids (Shanks *et al.*, 1997), and large scale production of secondary metabolites (Tikhomiroff and Jolicoeur, 2002). Cell cultures have higher rates of metabolism than intact differentiated plants because the initiation of cell growth in the culture leads to fast proliferation of cell mass and to a condensed biosynthetic cycle. This is the most important advantage of plant cell cultures as model systems for the study of the biosynthetic pathways, as secondary metabolite formation can take place within a short cultivation time, about 2 - 4 weeks (Dörmenburg and Knorr, 1995).

Although undifferentiated cell cultures have been studied, a large research interest has also been shown in the hairy root cultures (Bourgaud *et al.*, 2001). Hairy roots, a result of successful transformation of plant with *Agrobacterium rhizogenes*, is a possible source of secondary metabolites (Berzin *et al.*, 2000) because of the genetic and biochemical stability (Young-Am *et al.*, 2000), high level of differentiation as compared to cell



suspensions, amenability to genetic transformation (Shanks *et al.*, 1997) and rapid growth rate (Hamill and Lidgett, 1997; Shanks and Morgan, 1999). Hairy roots have been successfully applied in the production of some alkaloids (Tikhomiroff and Jolicoeur, 2002) such as  $\beta$ -carboline glucoid, ruine and serotonin (Misawa, 1994). However the only set-back is the lack of technology on how to scale up the hairy root culture to an industrial scale for a more economical process.

*Solenostemon scutellarioides* or locally name hati-hati has been known as a source of rosmarinic acid (Razzaque and Ellis, 1977) and also contains high levels of anthocyanins (Misawa, 1994). Rosmarinic acid possesses antioxidative properties (Tada *et al.*, 1996), antibacterial and anti-inflammatory effects (Petersen and Simmonds, 2003). It can be also used to prevent bronchial asthma, apasmogenic disorders, hepatotoxicity, ischaemic heart disease, and cancer disease (Al-Sereiti *et al.*, 1999) and can against human immunodeficiency virus types1 (HIV-1) (Mazunder *et al.*, 1997). Many studies on rosmarinic acid production by callus and cell suspension cultures of *Labiatae* family have been reported (Mizukami *et al.*, 1993; Su and Lei, 1993; Kintzios *et al.*, 1999). However, very limited studies have been carried out on medium optimization of callus induction and cell suspension culture. In addition, there has yet to be any report on *in vitro* propagation of multiple shoot, hairy root and normal root of *S. scutellarioides*. Therefore, the objectives of this research are:

1. To propagate *in vitro* cultures of *S. scutellarioides*.